

Lyme Neuroborreliosis with Long Meningeal Enhancement and Ischemic Stroke

Payam Dibaj^{a, b, c} Margret Schnegelsberg^c Renate Kröger^d

^aCenter for Rare Diseases Göttingen, University Medical Center Göttingen, Georg August University, Göttingen, Germany; ^bMax-Planck-Institute for Experimental Medicine, Göttingen, Germany; ^cCenter for Neurology, Asklepios Hospitals Schildautal, Seesen, Germany; ^dDepartment of Radiology, Asklepios Hospitals Schildautal, Seesen, Germany

A 61-year-old man with a 2-year history of prostate cancer presented with a slight hemiparesis on the left side that existed since 2 weeks. Additionally, the patient suffered from mild headache and decreased clinical condition. Neither fever nor signs of meningitis were observed in the clinical examination. Brain MRI revealed a corresponding restricted diffusion in the right internal capsule (Fig. 1a) and, accidentally, leptomeningeal enhancement after the injection of gadolinium at the level of the brainstem (Fig. 1b, c). Spinal MRI revealed similar meningeal enhancement over the whole spinal cord (Fig. 1d). The diagnosis was potentially leptomeningeal carcinomatosis, although it is of rare occurrence in patients with prostate cancer [1], but cytological analysis of the CSF showed predominant lymphocytic pleocytosis without malignant cells (178 cells/ μ L, protein 2.6 g/L, and lactate 4.2 mmol/L). In the CSF, CXCL13 was distinctly elevated (>500 pg/mL) [2, 3] and local antibody response to *Borrelia burgdorferi* was detected (IgG antibody index >17). Lyme neuroborreliosis with distinct meningeal enhancement was concluded as the most likely diagnosis. Nerve-root or men-

ingeal enhancement can rarely be observed in neuro-Lyme disease [4, 5]. Slight neuroborreliosis-associated cerebral vasculitis was considered to be the reason for the ischemic lesion [3, 6]. The hemiparesis of the patient subsequently improved, probably spontaneously. A treatment with ceftriaxone of 3 weeks' duration noticeably improved the clinical condition of the man and CSF findings. The case features two relatively rare clinical findings in neuroborreliosis: meningeal enhancement and ischemic stroke.

Acknowledgement

The authors are thankful to Prof. Sanjeev Sharma for careful reading and correction of the manuscript.

Statement of Ethics

The study was approved by the Ethics Committee of Asklepios clinics. Written informed consent was obtained from the patient for publication of the case including images.



Fig. 1. Diffusion-weighted imaging shows restricted diffusion in the right internal capsule (a), meningeal enhancement at the level of the brainstem (marked by an arrow) in the T1-weighted imaging after injection of gadolinium (b), axial plane at the level of the brainstem showing the leptomeningeal enhancement (arrow) (c), meningeal enhancement over the whole spinal cord (indicated between the two arrows) (d). Further imaging of the brain did not show signs of vasculitis or macroangiopathy.

Conflict of Interest Statement

There are no potential conflicts of interest with respect to the research, publication of this article, and/or financial completion.

Funding Sources

The authors did not receive any funding.

References

- 1 Carroll RD, Leigh EC, Curtis Z, Thorpe A, Ballengee J, Pacioles T. A case of leptomeningeal carcinomatosis from aggressive metastatic prostate cancer. *Case Rep Oncol.* 2019 Apr 10;12(1):311–6.
- 2 Schmidt C, Plate A, Angele B, Pfister HW, Wick M, Koedel U, et al. A prospective study on the role of CXCL13 in Lyme neuroborreliosis. *Neurology.* 2011 Mar 22;76(12):1051–8.
- 3 Garcia-Monco JC, Benach JL. Lyme neuroborreliosis. *Ann Neurol.* 2019 Jan;37(6):691–702.
- 4 Niksefat M, Albashiti B, Burke D, Moshayedi P, Patira R, Knepper L. Extensive meningeal enhancement in acute central nervous system Lyme: case series and review of literature. *J Clin Neurosci.* 2019 Jun;64:25–7.
- 5 Agarwal R, Sze G. Neuro-lyme disease: MR imaging findings. *Radiology.* 2009 Oct; 253(1):167–73.
- 6 Back T, Grünig S, Winter Y, Bodechtel U, Guthke K, Khati D, et al. Neuroborreliosis-associated cerebral vasculitis: long-term outcome and health-related quality of life. *J Neurol.* 2013 Jun;260(6):1569–75.

Author Contributions

P.D.: study concept and design, acquisition of data, analysis and interpretation, and writing of the manuscript; M.S.: analysis and interpretation, and critical revision of the manuscript; and R.K.: acquisition of data, analysis and interpretation, and critical revision of the manuscript.